



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Interactions between Autophagy and the Unfolded Protein Response: Implications for Inflammatory Bowel Disease

**Citation for published version:**

Hooper, K, Barlow, P, Henderson, P & Stevens, C 2018, 'Interactions between Autophagy and the Unfolded Protein Response: Implications for Inflammatory Bowel Disease', *Inflammatory Bowel Diseases*.  
<https://doi.org/10.1093/ibd/izy380>

**Digital Object Identifier (DOI):**

[10.1093/ibd/izy380](https://doi.org/10.1093/ibd/izy380)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Inflammatory Bowel Diseases

**Publisher Rights Statement:**

This is the author's peer-reviewed manuscript as accepted for publication

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



1 Interactions between Autophagy and the Unfolded Protein

2 Response: Implications for Inflammatory Bowel Disease

3  
4 Kirsty M. Hooper<sup>1</sup>, Peter G. Barlow<sup>1</sup>, Paul Henderson<sup>2, 3¶</sup> and Craig Stevens<sup>1¶\*</sup>.

5  
6 1. School of Applied Sciences, Edinburgh Napier University, Sighthill Campus, Sighthill Court,  
7 Edinburgh, EH11 4BN.

8 2. Child Life and Health, University of Edinburgh, Edinburgh, EH9 1UW.

9 3. Department of Paediatric Gastroenterology and Nutrition, Royal Hospital for Sick Children,  
10 Edinburgh, EH9 1LF.

11 ¶Joint senior authors

12  
13 Short title: Autophagy and UPR in IBD

14  
15 \*Address for Correspondence:

16 Dr Craig Stevens

17 School of Applied Sciences, Edinburgh Napier University, Sighthill Campus, Sighthill Court,  
18 Edinburgh, EH11 4BN.

19 Email: [C.Stevens@napier.ac.uk](mailto:C.Stevens@napier.ac.uk)

20 Tel: 0044 131 455 2930

21

22

## Abstract

Inflammatory Bowel Disease (IBD), which includes Crohn's disease (CD) and ulcerative colitis, is characterised by chronic inflammation of the gastrointestinal tract. Aetiology involves a combination of genetic and environmental factors resulting in abnormal immune responses to intestinal microbiota. Genetic studies have strongly linked genes involved in autophagy to CD, and genes involved in the unfolded protein response (UPR) to IBD. The UPR is triggered in response to accumulation of misfolded proteins in the endoplasmic reticulum (ER) and autophagy plays a key role to relieve ER-stress and restore homeostasis. This review summarises the known interactions between autophagy and the UPR and discusses the impact of these converging pathways on IBD pathogenesis. With a paucity of effective long-term treatments for IBD, targeting of synergistic pathways may provide novel and more effective therapeutic options.

**Keywords:** IBD, autophagy, unfolded protein response, ER stress.

## 37 Introduction

38 Inflammatory Bowel Disease (IBD) is a group of inflammatory diseases that includes Crohn's  
39 disease (CD), ulcerative colitis (UC) and IBD unclassified (IBDU). The incidence rate for IBD is  
40 approximately 50-200 in 100,000 persons per year in Western countries [1] and following  
41 diagnosis the natural history of the condition is characterized by periods of relapse and  
42 remission, with symptoms commonly including abdominal pain, chronic diarrhoea, weight  
43 loss and lethargy [2]. CD is distinguished from UC due to the presence of submucosal or  
44 transmural inflammation and macroscopic changes that often occur in a non-contiguous  
45 pattern anywhere within the digestive tract [1]. UC is localised to the colon and inflammation  
46 is limited to the mucosa and epithelial lining of the gastrointestinal (GI) tract [2]. Patients can  
47 be diagnosed with IBDU when a conclusive distinction between CD and UC cannot be made,  
48 although this may well represent a distinct sub-type. At present there is no cure for IBD and  
49 medications such as corticosteroids, aminosalicylates, immunomodulators and biological  
50 agents are aimed at inducing and maintaining remission of disease by modifying inflammatory  
51 processes [3].

52 The aetiopathology of IBD is multifactorial in nature, with genetic predisposition,  
53 environmental triggers (e.g. smoking, appendicectomy, diet, pollution, antibiotics and stress)  
54 and a dysregulated mucosal immune response contributing to disease [4]. Examination of the  
55 gut microbiome has revealed that IBD is associated with microbial dysbiosis, including an  
56 expansion of facultative anaerobic bacteria of the family Enterobacteriaceae [5]. Several  
57 potentially causative agents have been identified, most notably *Escherichia coli* strains with  
58 an adherent and invasive phenotype (AIEC) are associated with ileal mucosa in CD [6].  
59 Genome wide association studies (GWAS) have identified 240 IBD susceptibility loci to date

[7], and have confirmed association with previously recognised susceptibility genes including *Nucleotide-binding oligomerisation domain-containing protein 2 (NOD2)*. GWAS have also identified the strong association of CD with genes involved in the autophagy pathway, including *autophagy-related protein (ATG)16L1*, *Immunity-related GTPase family M protein (IRGM)* and *leucine rich repeat kinase 2 (LRRK2)* [8]. The strong association of IBD with endoplasmic reticulum (ER) stress/Unfolded protein response (UPR) genes including *x-box-binding protein 1 (XBP1)* [9] and genes involved in intestinal barrier function such as *MUC2* [10] and *Anterior gradient 2 (AGR2)* [11] have been detected by gene targeted approaches. Together, these genetic studies have led to increased research exploring links between autophagy and ER stress/UPR dysregulation and IBD pathogenesis.

## Autophagy

Autophagy is an intracellular process that plays an important housekeeping role by degrading excessive, damaged or aged proteins and organelles to maintain cellular homeostasis [12]. Basal autophagy is tightly regulated by the coordinated activity of autophagy-related (ATG) proteins [13] and constitutes an important survival mechanism induced in response to multiple stress conditions such as nutrient deprivation, hypoxia, DNA damage or intracellular pathogens [12]. There are three main types of autophagy in mammalian cells; macroautophagy (herein referred to as autophagy), microautophagy and chaperone-mediated autophagy [12].

When autophagy is initiated a double membrane vesicle is formed (the autophagosome) around the cargo to be degraded (**Figure 1**). The mature autophagosome then fuses with a lysosome to form an autophagolysosome, in which lysosomal enzymes degrade the inner

membrane and cargo and the resulting macromolecules are released into the cytosol for recycling (**Figure 1**).

Selective types of autophagy also exist, including autophagy of microorganisms (xenophagy) and autophagy of the ER membrane (ER-phagy), which use specific receptors and adaptor proteins to link the cargo to the autophagy machinery [14]. For example, Sequestosome 1/p62-like receptors (SLRs) target cytosolic pathogens and other cargo to initiate autophagy [15]. SLRs function by binding to the small regulatory protein ubiquitin on the surface of cargo [16–18] and subsequently associate with the autophagy machinery via a binding motif called the LC3-interacting region (LIR) [19]. Adaptor proteins, such as autophagy-linked FYVE protein (ALFY), can also bind ubiquitinated pathogens via p62 to promote association with the autophagy machinery [20]. To date, five main types of SLR have been described; sequestosome 1/p62, optineurin [18], NBR1 (Neighbor of BRCA1 gene 1) [21], NDP52 (Nuclear Domain 10 Protein 52) [17] and the NDP52-like receptor calcoco3 (Calcium-binding and coiled-coil domain-containing protein 3) [22], and specific cargo receptors are important for distinct types of selective autophagy. For example, a recent study has shown that the non-canonical cargo receptor cell-cycle progression gene 1 (CCPG1) is essential for ER-phagy [23], while another study demonstrated an integral role for optineurin in the maintenance of ER homeostasis by assisting the removal of hyper-activated UPR kinases [24].

## Autophagy and CD

Autophagy affects many essential cellular processes and dysregulation of autophagy has been linked to a multitude of human diseases [25]. Autophagy plays an important role in both innate and adaptive immune signalling pathways and loss of immune regulation is a key event

leading to the chronic inflammation observed in CD [26]. Impaired autophagy responses have been observed in a range of cell types derived from CD patients including the specialized intestinal epithelial cells (IECs) Paneth cells and goblet cells, and leukocytes, such as macrophages and dendritic cells [27].

Functional studies have linked impaired autophagy to CD-associated genetic variants in *NOD2*, *ATG16L1*, *IRGM* and *LRRK2*. The single nucleotide polymorphism (SNP) in *ATG16L1* causes a single amino acid change from threonine to alanine at position 300 (*T300A*) [28], which is associated with Paneth cell and goblet cell dysfunction, and significantly impairs autophagic clearance of pathogens [29–32]. *IRGM* is required for the initiation of xenophagy and the clearance of intracellular organisms such as *Mycobacterium tuberculosis* [33] and dysregulation of *IRGM* expression compromises the control of intracellular replication of CD-associated adherent invasive *Escherichia coli* (AIEC) by autophagy [34]. *LRRK2* expression is increased in colonic biopsy specimens from patients with CD [35] and functionally *LRRK2* can enhance NFκB-dependent transcription, while small interfering RNA [siRNA] knockdown of *LRRK2* interferes with bacterial killing [35].

*NOD2* is a member of the Nod-like receptor (NLR) family of pattern recognition receptors (PRR) and recognises a component of the bacterial cell wall muramyl dipeptide (MDP) to induce innate immune responses [36]. CD-associated *NOD2* SNPs (R702W, G908R and L1007fs) affect the leucine rich repeat domain disrupting interaction with MDP and abrogating immune responses initiated by this receptor [37]. The immunoregulatory properties of *NOD2* have also been linked to autophagy, and CD susceptibility is heightened when *ATG16L1* and *NOD2* variants present in combination, causing synergistic genetic epistasis [38,39]. A direct functional interaction between these proteins has been

determined; NOD2 was shown to recruit ATG16L1 to the plasma membrane to initiate autophagy at the sites of bacterial entry [40], and in a separate study IRGM was shown to regulate the formation of a complex containing NOD2 and ATG16L1 that is necessary for the induction of xenophagy [41]. The interaction of IRGM with NOD2 also stimulates phosphorylation cascades involving AMPK, ULK1 and Beclin1 that regulate autophagy initiation complexes [41]. Cells harbouring CD-associated *NOD2* variants and/or the *ATG16L1 T300A* variant exhibit a number of disrupted functions linked to autophagy including reduced production of antimicrobial peptides, enhanced pro-inflammatory responses and aberrant activation of adaptive immune responses [40,42–44].

Significantly, abnormalities in the secretory capacity of Paneth cells are observed in mice deficient for ATG16L1 [30,45,46], NOD2 [47,48], IRGM [49] and LRRK2 [50] indicating that autophagy plays an essential and specific role in Paneth cell function. Despite the significant effects on Paneth cell function, mouse strains developed for deficiency in functional ATG16L1 do not exhibit spontaneous intestinal inflammation [29–31]. In contrast, a mouse strain with targeted deletion of *ATG16L1* in IECs developed a spontaneous transmural ileitis similar to ileal CD [24]. Furthermore, targeted deletion of *ATG16L* in haematopoietic cells can enhance susceptibility to DSS-induced acute intestinal injury in mice [51] and ATG16L1 deficiency in myeloid cells in a mouse strain led to disrupted macrophage function and bacterial clearance [52]. Murine models with non-functional NOD2 do not develop spontaneous colitis [53], however a *NOD2* mutation similar to the L1007fs mutation increased susceptibility to DSS-induced colitis in mice [54]. *Irgm1*-deficient mice also exhibit abnormalities in Paneth cells, accompanied by increased susceptibility to inflammation in the colon and ileum [49]. Finally, *LRRK2* deficiency confers enhanced susceptibility to experimental colitis in mice, which was



associated with enhanced nuclear localisation of the transcription factor nuclear factor of activated T cells (NFAT1), important for regulating innate immune responses [55].

## ER-stress and UPR signalling

ER stress results from accumulation of unfolded and misfolded protein in the ER, and the UPR is activated to resolve ER stress and restore homeostasis. The UPR inhibits protein synthesis, promotes protein re-folding, and induces degradation of unfolded and misfolded proteins through ER-associated protein degradation (ERAD) and autophagy (**Figure 2**). If these survival mechanisms are unsuccessful, the UPR can induce apoptosis [56]. The major regulators of the UPR are the ER-membrane resident proteins PERK (protein kinase RNA-like endoplasmic reticulum kinase), inositol-requiring transmembrane kinase endonuclease 1 (IRE1) and activated transcription factor (ATF)6. When inactive these proteins are bound to binding immunoglobulin protein (BiP), also known as glucose regulated protein 78 (GRP78) [57]. During ER stress, BiP binds to misfolded proteins in the ER and dissociates from the ER-membrane resident proteins to allow their transition to an active state [57] (**Figure 2**).

When active, PERK phosphorylates elongation initiation factor  $2\alpha$  (EIF2 $\alpha$ ), to inhibit general protein synthesis [58] and specifically up-regulates ATF4 [59]. ATF4 in turn transcriptionally up-regulates several other UPR genes including CCAAT/enhancer-binding protein (C/EBP) homologous protein (CHOP) [60,61] (**Figure 2**). CHOP is also a transcription factor that regulates several other UPR genes, and under conditions of prolonged ER stress can promote apoptosis [60,61].

IRE1 exists in two forms: IRE1 $\alpha$  that is ubiquitously expressed and IRE1 $\beta$  that is only expressed in the GI tract and lung epithelial cells [62]. During ER stress, IRE1 is activated through

dimerization and auto-phosphorylation [63,64]. The IRE1 $\alpha$  RNase domain is essential for creating transcriptionally activate *XPB1* messenger RNA (mRNA) via splicing, which acts as a transactivator of UPR genes [65–68] (**Figure 2**) . IRE1 endoribonuclease activity also facilitates degradation of specific mRNA in a process known as RIDD (regulated IRE1-dependent decay) [69].

ATF6 translocates to the Golgi apparatus once released from its complex with BiP [70]. This allows cleavage by site 1 and site 2 proteases (S1P and S2P), which releases the transcriptionally active cytoplasmic domain of ATF6 (ATF6-N) that induces UPR-associated genes [71–73] (**Figure 2**). Among the ATF6 upregulated genes are *CHOP* and *XPB1* [74].

## ER-stress, UPR and intestinal inflammation

Genetic studies have identified several ER-stress/UPR genes associated with IBD [75]. Moreover, ER-stress levels are increased in ileal and colonic biopsies from CD patients, with higher than normal levels of BiP, chaperone protein Gp96, and spliced *XPB1* observed [9,76–78] (**Table 1**). Several studies have focused on IRE1-*XPB1* signalling in murine models. In mice with targeted deletion of *XPB1* in intestinal epithelial cells (IECs) (*XPB1* <sup>$\Delta$ IEC</sup> mice), spontaneous inflammation of the small intestine, increased susceptibility to DSS-induced colitis and elevated levels of ER stress were observed [9] (**Table 1**). Furthermore, in *XPB1* <sup>$\Delta$ IEC</sup> mice increased levels of apoptosis were observed along with reduced goblet cell and Paneth cell numbers, leading to decreased production of host defence peptides and higher susceptibility to *Listeria monocytogenes* infection [9] (**Table 1**). *XPB1* has also been shown to suppress experimental colitis-associated cancer [79], and is essential for efficient TLR-mediated pro-

inflammatory responses to infection in macrophages [80]. These studies support *XBP1* as a key component of the protective function of IECs and macrophages.

Although the UPR acts to maintain ER-homeostasis, hyper-activation of certain UPR components can create a pro-inflammatory state. In *XBP1* <sup>$\Delta$ IEC</sup> mice increased activation of IRE1 $\alpha$ , causes hyper activation of NF $\kappa$ B, and spontaneous inflammation [45] (**Table 1**). IRE1 $\beta$  knock-out mice have enhanced sensitivity to DSS-induced colitis [81] and exhibit goblet cell abnormalities with exaggerated MUC2 accumulation (**Table 1**). In contrast, IRE1 $\alpha$  knock-out mice have normal goblet cells [82]. In murine Paneth cells, IRE1 $\alpha$  and IRE1 $\beta$  have distinct roles with hyper activation of IRE1 $\alpha$  driving CD-like ileitis, and IRE1 $\beta$  having a protective role [24].

Association of aberrant PERK-EIF2 $\alpha$  and ATF6 pathways with intestinal inflammation have also been identified. A mouse model expressing non-phosphorylatable EIF2 $\alpha$  in IECs resulted in functional abnormalities in Paneth cells and increased susceptibility to *Salmonella* infection and DSS-induced colitis [83] (**Table 1**). ATF6 $\alpha$  deficient mice exhibit increased ER stress as indicated by elevated levels of BiP, ATF4, CHOP and spliced *XBP1*, which result in enhanced sensitivity to DSS-induced colitis [84] (**Table 1**). Additionally, hypomorphic mutation in *membrane-bound transcription factor peptidase S1P-encoding gene (Mbtps1)*, which encodes the S1P responsible for cleavage of ATF6, causes enhanced susceptibility to DSS-induced colitis [85]. Although there is less evidence to support a role for PERK-EIF2 $\alpha$  and ATF6 pathways in IBD pathogenesis, their importance for ER stress responses in the intestinal epithelium is clear.

## ER-stress and intestinal barrier function

In the intestinal epithelium, cells that naturally secrete large amounts of protein, such as Paneth cells and goblet cells, are more susceptible to ER-stress and therefore rely heavily on the UPR to maintain homeostasis. MUC2 is the major component of mucin that is produced in goblet cells and secreted into the intestinal lumen. *Winnie* mice are characterised by a missense mutation in *MUC2*, which causes abnormalities in goblet cells, leading to aberrant mucous production and spontaneous colitis, and association with *MUC2* variants has been identified in IBD patients [86]. *Winnie* mice also exhibit severe ER stress in goblet cells [10], which causes up to four-fold increase in activated dendritic cells in the colonic lamina propria, and aberrant adaptive immune responses associated with interleukin (IL)-23/Th17 [87]. Goblet cell abnormalities are also apparent in mice deficient in UPR transcription factor OASIS, which causes increased ER stress and susceptibility to DSS-induced colitis [88,89]. AGR2 is an ER resident protein highly expressed in goblet and Paneth cells and regulates the formation of disulphide bonds in mature proteins. *AGR2*<sup>-/-</sup> mice exhibit a decreased number of goblet cells and MUC2 production, Paneth cell abnormalities, elevated ER-stress and spontaneous colitis [90]. Notably, AGR2 is decreased in patients with CD and UC [11]. These studies highlight the key role of intestinal secretory cells and breakdown of intestinal barrier function in IBD pathogenesis.

## Functional intersection between autophagy and the UPR

The UPR and autophagy are intimately linked processes. In a range of Intestinal epithelial cells, chemical ER-stress inducers activate autophagy, modulated by enhanced expression of CHOP and stimulation of the IRE1 $\alpha$  pathway [91]. In endothelial cells, IRE1 $\alpha$ -dependent splicing of *XBP1* mRNA activated autophagy via up-regulation of *Beclin-1*, which is a major regulator of

the autophagy pathway [92] (**Figure 3**). Contrary to expectation, *XPB1* deletion in a familial amyotrophic lateral sclerosis mouse model increased autophagy, which enhanced clearance of accumulated toxic superoxide dismutase-1 (SOD1) aggregates [93]. It was suggested that in this scenario, autophagy is induced in a compensatory manner due to attenuated UPR.

The UPR and autophagy also intersect at the PERK-EIF2 $\alpha$ -ATF4 pathway [94–99]. In an *in vitro* model of osteosarcoma, PERK induced autophagy via mechanistic target of rapamycin (mTORC1) inhibition to promote survival in response to ER stress-conferred chemoresistance to apoptosis [95] (**Figure 3**). Additionally, PERK modulates autophagy via AMPK-dependent inhibition of mTORC1 in response to extracellular matrix (ECM) detachment in mammary epithelial cells (MECs) [94]. One of the main functional outcomes of PERK signalling is reduced protein synthesis. Inhibition of mTORC1 helps to promote this effect as mTORC1 controls synthesis of ~15-20% of protein within the cell [100]. Thus, via modulation of mTORC1, PERK signalling achieves dual outcomes; inhibition of protein synthesis and induction of autophagy to degrade misfolded proteins.

During amino acid deprivation, ATF4 and CHOP can bind specific C/EBP-ATF Response Elements (CAREs), also known as Amino Acid Response Elements (AAREs) and CHOP-Response Elements (CHOP-REs) to induce transcription of a wide range of autophagy genes [101] (**Figure 3**). In other studies, hypoxia or ECM detachment induced PERK-dependent autophagy due to autophagy gene up-regulation via ATF4 and CHOP [102–104]. This up-regulation of autophagy gene transcription by the UPR was shown to replenish autophagy proteins to promote survival during cellular stress [103].

ATF6 has also been implicated mechanistically in autophagy regulation. In response to cellular stress, interferon (IFN)- $\gamma$  activates the Ask1 (Apoptosis signal-regulating kinase 1)/MAPK (Mitogen-activated protein kinase) pathway, which phosphorylates ATF6 to allow its proteolytic activation [105]. ATF6 interaction with C/EBP- $\beta$  is essential for IFN- $\gamma$ -induced up-regulation of *DAPK1* (*death-associated protein kinase 1*), which can subsequently stimulate autophagy [106] (**Figure 3**). Mice lacking either ATF6 or Ask1 are highly susceptible to bacterial infection due to defective autophagy [105,106]. Furthermore, ATF6 recruitment of DAPK1 in response to ER stress enhanced xenophagy in human colonic biopsies and epithelial cells, which was attenuated in cells harbouring the *ATG16L1 T300A* SNP [107]. Additionally, activated ATF6 was shown to stimulate Akt (protein kinase B), which resulted in the inhibition of mTORC1 [108,109] (**Figure 3**).

In a recent study in MCF-7 human breast cancer cells, ER stress induced by the chemopreventative agent ursolic acid (UA) was associated with autophagy activation [99]. UA induced autophagy via MAPK1/3 signalling and subsequent promotion of PERK signalling, resulting in the inhibition of apoptosis. Furthermore, a study in human ovarian cancer cells showed interdependent activation of autophagy and the PERK-EIF2 $\alpha$  UPR pathway when treated with metformin, which causes energy starvation [98]. In these scenarios an unconventional relationship between autophagy and ER stress was uncovered, which remains to be mechanistically solved. Nonetheless, under these circumstances the interaction of the UPR and autophagy pathways has pro-survival outcomes.

## Convergence of autophagy, ER-stress and CD

In an attempt to relieve ER-stress the UPR can induce autophagy to degrade misfolded proteins, protein aggregates and damaged organelles [91,110–113]. Autophagy activity is increased in highly secretory Paneth cells [45] to counterbalance high levels of ER-stress [112], thus ER-stress is a significant risk in these cells when the UPR or autophagy is not functional. Consistent with this, in Paneth cells of CD patients harbouring *ATG16L1 T300A* risk alleles, BiP and pEIF2 $\alpha$  are highly expressed [46] (**Table 1**). Significantly, *ATG16L1;XBP1<sup>ΔIEC</sup>* mice develop similar phenotypic ileitis to *ATG16L1<sup>ΔIEC</sup>* mice, but earlier in life due to increased ER stress [24,45].

ERAD can regulate the degradation of IRE1 $\alpha$  to prevent accumulation of toxic IRE1 $\alpha$  aggregates, however persistent ER-stress will inhibit ERAD degradation of IRE1 $\alpha$  [24]. When this occurs, autophagy plays an important role in the clearance of supramolecular clusters of IRE1 $\alpha$  (**Figure 3**). In *ATG16L1<sup>ΔIEC</sup>* mice, development of spontaneous CD-like ileitis is associated with defective autophagy resulting in toxic accumulation of IRE1 $\alpha$  in Paneth cells [24] (**Table 1**). Furthermore, the selective autophagy receptor optineurin interacts with IRE1 $\alpha$ , and optineurin deficiency amplified the accumulation of IRE1 $\alpha$  [24]. In humans homozygous for *ATG16L1 T300A*, a similar accumulation of IRE1 $\alpha$  was observed in intestinal epithelial crypts [24] (**Table 1**). This has led to suggestion that the *ATG16L1 T300A* SNP may define a specific subtype of patients with CD, characterised by Paneth cell ER-stress [46]. This synergistic and compensatory relationship between the UPR and autophagy is affirmed by the presence of CD-associated SNPs in *ATG16L1* and *XBP1*.

A recent study has demonstrated a direct link between NOD1/2 and the IRE1 $\alpha$  pathway in the context of ER-stress-induced inflammation [114]. When active, IRE1 $\alpha$  stimulates the c-Jun N-terminal kinase (JNK) pathway and recruits TRAF2 (TNF receptor-associated factor 2) to the

ER membrane to trigger NF $\kappa$ B signalling [115,116] and autophagy induction [112,117,118] (Figure 3). In mouse and human cells, ER-stress induced by chemicals or infection with *Brucella abortus* and *Chlamydia muridarum* increased inflammation and IL-6 production [114]. This response was dependent on NOD1/2 and receptor-interacting serine/threonine-protein kinase 2 (RIPK2), but also on IRE1 $\alpha$  kinase activity and TRAF2-induced NF $\kappa$ B signalling [114]. This suggests there is a functional intersection between the IRE1 $\alpha$  pathway and NOD1/2 signalling, which is facilitated by TRAF2 (Figure 3).

Interestingly, an additional study has shown that ER-stress responses can be modulated by another innate immune sensor called stimulator of interferon genes (STING) in response to cyclic-di-AMP (c-di-AMP), a vita-PAMP (pathogen associated molecular pattern) present in live Gram-positive bacteria [119]. This process induces autophagy via inhibition of the major autophagy suppressor mTORC1 and localisation of STING to autophagosomes.

## Pharmacological induction of autophagy and the UPR

A recent review estimated IBD treatment costs of £720 million (\$940m) per year in the United Kingdom alone [120], with roughly a quarter of these costs directly attributed to drug treatments [121]. The efficacy of these drugs continues to come under scrutiny as response to treatment often diminishes over time, with a review of worldwide cohorts estimating that between 10–35% of CD patients required surgery within a year of diagnosis and up to 61% by 10 years [122]. In order to improve the efficacy of IBD treatment, optimization of existing clinical therapies and the development of novel therapeutics is required.

The convergence between autophagy and UPR pathways provides new opportunity for the treatment of IBD and the modulation of the UPR in combination with autophagy inducers is a



promising therapeutic strategy. There is evidence that inducing autophagy can have therapeutic benefits for the treatment of IBD [26] with several studies investigating the utility of autophagy inducers as adjuvant therapies. Rapamycin analogues, sirolimus and everolimus, inhibit mTORC1 to induce autophagy and are already approved for clinical use for post-transplantation (e.g. liver and renal) management. In IL-10-deficient mice, everolimus treatment alleviated spontaneous colitis and reduced CD4<sup>+</sup> T cells and IFN- $\gamma$  [123]. In a case study sirolimus improved symptoms and intestinal healing in a patient with severe refractory CD [124]. In another case study, symptoms were controlled for 18 months with everolimus treatment in a refractory UC patient [125]. Moreover, in a study of refractory paediatric IBD, sirolimus induced clinical remission in 45% of UC patients and 100% of CD patients; albeit the sample size was small [126]. Significantly, everolimus had comparable safety and tolerability as azathioprine when used to maintain steroid-induced remission in a cohort of adult CD patients [127]. As these mTORC1 inhibitors are already approved for clinical use, they have been investigated the most extensively, however there are a plethora of novel autophagy modulators that are currently being developed, characterised and patented for therapeutic use in a range of diseases including IBD [128,129].

Recent progress has also been made to identify specific chemical inducers of the UPR. A screen of 1,200 FDA-approved compounds carried out in *C.elegans* identified eight compounds that induced UPR responses, four of which specifically increased mitochondrial UPR [130]. The identified drugs included antirheumatic agents, antianginal calcium channel blockers; androgen receptor inhibitors used for cancer therapy and tetracycline antibiotics.

A well-characterised modulator of the UPR, tauroursodeoxycholic acid (TUDCA), that promotes protein refolding to reduce ER-stress, was shown to ameliorate DSS-induced colitis

in mice by decreasing ER-stress in IECs [84]. Furthermore, a selective inhibitor of eIF2 $\alpha$  dephosphorylation protects cells from ER-stress and ameliorates murine experimental colitis [131,132]. Supplementation with glutamine has also been suggested for the improvement of IBD treatment, as this amino acid was shown to dampen experimental colitis in rats by inhibiting ER-stress in colonic epithelial cells [133].

Drugs used to treat metabolic disorders have also been investigated for UPR inducing properties. The biguanides metformin and phenformin have been implicated in induction of the UPR and resolution of ER-stress via activation of AMPK, which subsequently stimulated IRE1 $\alpha$  and PERK pathways [98,134,135]. Inhibitors of dipeptidyl peptidase IV (DPP4), including gemigliptin, also prevented ER-stress-mediated apoptosis by promoting IRE1 $\alpha$  and PERK pathways [136]. Furthermore, agonists of the glucagon-like peptide-1 receptor, such as exenatide, relieved ER stress via up-regulation of *ATF4* expression [137]. Exogenous chemical chaperones have also been explored as a method to relieve ER stress by mimicking ER chaperones to promote protein transport and re-folding capacity [138].

Although several studies have demonstrated beneficial effects of enhancing UPR function for intestinal homeostasis, future investigations should proceed with caution. For example, hyper-activation of the UPR kinase IRE1 $\alpha$  can exacerbate intestinal inflammation, as seen in patients with *ATG16L1* and *NOD2* mutations, therefore, in certain circumstances pharmacological inhibition of UPR receptors would be a more effective strategy [24,45,114]

Of particular interest, the selective autophagy cargo receptor optineurin forms a critical link between ER-stress resolution and autophagy due to its role in the degradation of IRE1 $\alpha$  aggregates [24], and another recently identified autophagy cargo receptor that is integral for resolution of ER-stress, CCPG1, mediates ER-phagy to remove damaged ER membranes [23]. Understanding the biology and functions of adaptors such as optineurin and CCPG1 may identify novel druggable targets and expedite development of the next generation of therapeutics aimed at modulation of the UPR in combination with autophagy.

## Discussion

The complexity of IBD is evident from the large number of risk loci identified by genetic studies, and the diverse health profile of patients that are affected. Mouse models of IBD cannot emulate the human disease, however they are useful tools to explore how specific gene mutations influence inflammation. Interestingly, as highlighted in **(Table 1)** the majority of mouse models mimicking IBD-associated genetic risk do not develop spontaneous inflammation, but rather they are sensitised to DSS-induced colitis, which acts by damaging the epithelium and increasing intestinal permeability. The intestinal epithelium has important immunoregulatory functions and controls the equilibrium between tolerance and immunity to non-self-antigens [139]. As such breakdown of intestinal epithelial barrier function and concomitant interaction with environmental factors in the lumen is a trigger for inflammation. The intestinal lumen comprises a multitude of potential triggers including the microbiota, dietary antigens, and luminal antigens. Additional triggers may be host-derived factors that are released into the lumen as the intestinal epithelial barrier breaks down. These so-called Damage-Associated Molecular Patterns (DAMPS) include intracellular proteins, such as high-mobility group box 1 (HMGB1), heat-shock proteins and components derived from

the extracellular matrix. Examples of non-protein DAMPs include genomic DNA, mitochondrial DNA, RNA, uric acid and ATP [140,141]. Not surprisingly, there is considerable interest in developing novel therapeutic strategies aimed at re-establishing intestinal barrier function [142] and modulation of DAMPs for the treatment of IBD [140].

Dysbiosis of the gut microbiome is strongly implicated in the pathogenesis of CD [143], and it has been suggested that microbial dysbiosis may be an environmental trigger. A recent study by Tschurtschenthaler and colleagues [24] addressed this question. Although microbial dysbiosis was present in the ileum of *Atg16l1;Xbp1*<sup>ΔIEC</sup> mice, such structural alteration of the microbiota did not trigger ileitis but, rather, aggravated DSS-induced colitis [24]. In order to understand the role of the environment in disease, determining the relative contribution of genetics and a detailed characterization of environmental triggers is required.

Greater understanding of the genetic factors that underlie CD pathogenesis are leading to improvements in treatment. Development of personalised therapies may be achieved via genotyping for key SNPs in genes involved in both the autophagy and UPR pathways. IBD drugs already established in the clinic have been shown to exert their effects, at least in-part, through the modulation of autophagy [26] or the UPR, and establishing patient genotypes may help predict response. For example, recent studies have identified an association between *ATG16L1 T300A* SNP and an enhanced therapeutic effect of thiopurines [144] and anti-TNF-α therapy [145]. Interestingly, the immunoregulatory effects of these drugs were associated with autophagy stimulation [144,146,147] and the *T300A* genotype has been associated with a subset of patients that exhibit deficiencies in both the UPR and autophagy [46]. Furthermore, CD patients harbouring *NOD2* mutations associate with better clinical outcomes in response to thiopurines, whereas CD patients with wild-type *NOD2* respond

412 better to steroids and anti-TNF therapy [148]. Due to the genetic complexity of IBD and  
 413 epistasis between genes, it is imperative that multiple genes are analysed for the purpose of  
 414 patient stratification. For example, a recent study identified a 32-gene transcriptomic  
 415 signature in lymphoblastoid cells that was able to predict lack of response to thiopurines, with  
 416 aberrant cell cycle control, DNA mismatched repair and RAC1-dependent mechanisms  
 417 implicated in thiopurine resistance [149]. Furthermore, it is increasingly clear that epigenetic,  
 418 microRNA and immune cell signatures among others will have a significant role to play in  
 419 predicting disease susceptibility and response to therapy [150–152].

420 With regards to the intestinal microbiota, a recent study has characterised microbial  
 421 signatures for the diagnosis of IBD that were highly sensitive and could differentiate CD  
 422 patients from healthy controls and UC patients. This study highlights the potential for using  
 423 the intestinal microbiota as a micro-biomarker [153]. Importantly, as many drugs need to be  
 424 metabolised and de-toxified by the gut microbiota, this approach could also have application  
 425 in predicting response to therapy. Given that dysregulation of autophagy and ER-stress can  
 426 affect the intestinal microbial environment, analysis of microbial signatures may help to  
 427 determine if a patient would benefit from drugs that modulate the autophagy or UPR  
 428 pathways.

429 To conclude, the ER-stress/UPR and autophagy pathways play a vital role in the maintenance  
 430 of intestinal homeostasis and breakdown of these converging pathways has been implicated  
 431 in persistent intestinal infections, chronic inflammation and dysregulated immune responses  
 432 observed in IBD. Therefore, strategies aimed at modulating these pathways simultaneously  
 433 may prove to be an effective therapeutic option.

## 434 Funding

435 This work was supported by a Crohn's in Childhood Research Association (CICRA) PhD  
436 studentship to KMH. PH is supported by a NHS Research Scotland Career Researcher  
437 Fellowship.

## Figure Legends

### Figure 1: Autophagy pathway and autophagosome biogenesis

During the initial stages of autophagy, the isolation membrane forms a double membrane vesicle (the autophagosome) around the cargo to be degraded. ULK complex (ULK1-ULK2-ATG13-FIP200-ATG101) and Beclin 1 (Vps34-Vps150-Beclin1) complex, through interaction with ATG14, recruit autophagy proteins and complexes to the autophagosome membrane. ATG12 is conjugated to ATG5 and forms a complex with ATG16L1 (ATG16L1 complex). The ATG16L1 complex is proposed to specify the site of LC3 lipidation for autophagosome formation. LC3 is conjugated to PE to form lipidated LC3-II and is associated with the autophagosome outer membrane. Upon autophagosome closure, LC3 localises to the inner membrane and other autophagy proteins and complexes dissociate for recycling. The mature autophagosome then fuses with a lysosome to form an autophagolysosome, in which cargo are degraded by lysosomal enzymes and subunits are recycled.

### Figure 2: The unfolded protein response

BiP chaperone protein binds unfolded/misfolded proteins in the ER and dissociates from transmembrane receptors upon accumulation of the toxic proteins. The transmembrane receptors PERK, IRE1 $\alpha$  and ATF6 become activated. PERK phosphorylates EIF2 $\alpha$ , which downregulates global translation but specifically upregulates ATF4 and CHOP that upregulate UPR-associated genes. IRE1 $\alpha$  splices XBP1 to its active form and ATF6 is cleaved by S1P and S2P to active ATF6-N, which both translocate to the nucleus to upregulate UPR-associated genes. The main function of these UPR-associated genes is to increase protein refolding,

inhibit synthesis of new protein and degrade unfolded/misfolded proteins through autophagy and ERAD.

### Figure 3: Intersection between autophagy and the unfolded protein response

ER stress activates transmembrane receptors PERK, IRE1 $\alpha$  and ATF6. PERK phosphorylates EIF2 $\alpha$ , which specifically upregulates ATF4 and CHOP that bind AAREs and CHOP-Res to upregulate autophagy genes. PERK also induces autophagy via mTORC1 inhibition. IRE1 $\alpha$  splices XBP1 to its active form, which up-regulates *Beclin-1*. IRE1 $\alpha$  endonuclease activity activates the JNK pathway, which induces autophagy via TRAF2, NOD2 and NF $\kappa$ B. Enhanced autophagy degrades accumulated IRE1 $\alpha$  clusters. Active ATF6-N induces autophagy via mTORC1 inhibition and binds C/EBP- $\beta$  to up-regulate *DAPK1*.

### Table 1: Murine models of intestinal inflammation

Links between autophagy, ER-stress/UPR and experimental colitis/intestinal inflammation and IBD.






Figure 1




Figure 2




Figure 3

Autophagy/ UPR pathway	Murine models of intestinal inflammation	IBD patients
ATG16L1	<ul style="list-style-type: none"> <li>ATG16L1 deficiency caused enhanced susceptibility to experimental colitis, Paneth cell and Goblet cell dysfunction, disrupted macrophage function and significantly impairs xenophagy [29-32, 51, 52]</li> <li>ATG16L1 deletion in IECs induced spontaneous transmural ileitis [24]</li> </ul>	<i>ATG16L1 T300A</i> CD-associated SNP [28]
NOD2	NOD2 mutation causes enhanced susceptibility to DSS-induced colitis [54] and causes Paneth cell dysfunction [47, 48]	<i>NOD2</i> CD-associated SNPs (R702W, G908R and L1007fs) [37]
IRGM	<i>Irgm1</i> deficiency causes abnormalities in Paneth cells and increased susceptibility to inflammation in the colon and ileum [49]	<i>IRGM</i> CD-associated SNP [8]
LRRK4	LRRK2 deficiency confers enhanced susceptibility to experimental colitis in mice [55] and Paneth cell abnormalities [50]	<i>LRRK4</i> CD-associated SNP [8]
IRE1 $\alpha$ -XBP1	<ul style="list-style-type: none"> <li><i>XBP1</i> deletion causes spontaneous intestinal inflammation, abnormal Paneth and goblet cell function and increased infection [9]</li> <li><i>XBP1</i> deletion causes overactivation of IRE1<math>\alpha</math> and NF<math>\kappa</math>B [45]</li> <li><i>ATG16L1</i> deletion causes accumulation of IRE1<math>\alpha</math> in Paneth cells resulting in CD-like ileitis [24]</li> </ul>	<ul style="list-style-type: none"> <li><i>XBP1</i> CD-associated SNP [9]</li> <li>Increased levels of spliced <i>XBP1</i>, BiP and Gp96 in CD [9, 76-78]</li> <li><i>T300A</i> SNP causes accumulation of IRE1<math>\alpha</math> in intestinal crypts [24]</li> </ul>
IRE1 $\beta$	IRE1 $\beta$ deletion causes enhanced sensitivity to DSS-colitis [81], goblet cell abnormalities and MUC2 accumulation [24]	
PERK-EIF2 $\alpha$	Non-phosphorylatable EIF2 $\alpha$ caused Paneth cell abnormalities, enhanced DSS-colitis susceptibility and increased <i>Salmonella</i> infection [83]	Increased p-EIF2 $\alpha$ and BiP in CD patients with <i>T300A</i> SNP [46]
ATF6	<ul style="list-style-type: none"> <li>ATF6 deletion enhanced DSS-colitis susceptibility [84]</li> <li>Mutation in <i>Mbtps1</i> (encodes S1P) causes enhanced DSS-colitis susceptibility [85]</li> </ul>	
AGR2	AGR2 deletion causes decreased Goblet cells and MUC2 production, Paneth cell abnormalities, elevated ER-stress and spontaneous colitis [90]	<ul style="list-style-type: none"> <li><i>AGR2</i> CD-associated SNP [11]</li> <li><i>AGR2</i> decreased in IBD [11]</li> </ul>

## 480 Table 1

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- 1 Gasparetto M, Guariso G. Highlights in IBD Epidemiology and Its Natural History in the Paediatric Age. *Gastroenterol Res Pr* 2013;**2013**:829040. doi:10.1155/2013/829040
- 2 Fakhoury M, Negrulj R, Mooranian A, *et al.* Inflammatory bowel disease: clinical aspects and treatments. *J Inflamm Res* 2014;**7**:113–20. doi:10.2147/JIR.S65979
- 3 Neurath MF. Current and emerging therapeutic targets for IBD. *Nat Rev Gastroenterol Hepatol* 2017;**14**:nrgastro.2016.208. doi:10.1038/nrgastro.2016.208
- 4 Boyapati R, Satsangi J, Ho GT. Pathogenesis of Crohn's disease. *F1000Prime Rep* 2015;**7**:44. doi:10.12703/p7-44
- 5 Zuo T, Ng SC. The Gut Microbiota in the Pathogenesis and Therapeutics of Inflammatory Bowel Disease. *Front Microbiol* 2018;**9**:2247. doi:10.3389/fmicb.2018.02247
- 6 Darfeuille-Michaud A, Boudeau J, Bulois P, *et al.* High prevalence of adherent-invasive *Escherichia coli* associated with ileal mucosa in Crohn's disease. *Gastroenterology* 2004;**127**:412–21. doi:10.1053/j.gastro.2004.04.061
- 7 Lange KM de, Moutsianas L, Lee JC, *et al.* Genome-wide association study implicates immune activation of multiple integrin genes in inflammatory bowel disease. *Nat Genet* 2017;**49**:256–61. doi:10.1038/ng.3760
- 8 Franke A, McGovern DP, Barrett JC, *et al.* Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nat Genet* 2010;**42**:1118–25. doi:10.1038/ng.717
- 9 Kaser A, Lee A-H, Franke A, *et al.* XBP1 Links ER Stress to Intestinal Inflammation and Confers Genetic Risk for Human Inflammatory Bowel Disease. *Cell* 2008;**134**:743–56. doi:10.1016/j.cell.2008.07.021
- 10 Heazlewood CK, Cook MC, Eri R, *et al.* Aberrant Mucin Assembly in Mice Causes Endoplasmic Reticulum Stress and Spontaneous Inflammation Resembling Ulcerative Colitis. *PLoS Med* 2008;**5**. doi:10.1371/journal.pmed.0050054
- 11 Zheng W, Rosenstiel P, Huse K, *et al.* Evaluation of AGR2 and AGR3 as candidate genes for inflammatory bowel disease. *Genes Immun* 2006;**7**:11–8. doi:10.1038/sj.gene.6364263
- 12 Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy. *Nat Rev Mol Cell Biol* 2018;**19**:349. doi:10.1038/s41580-018-0003-4
- 13 Lamb CA, Yoshimori T, Tooze SA. The autophagosome: origins unknown, biogenesis complex. *Nat Rev Mol Cell Biol* 2013;**14**:759–74. doi:10.1038/nrm3696
- 14 Zaffagnini G, Martens S. Mechanisms of Selective Autophagy. *J Mol Biol* 2016;**428**:1714–24. doi:10.1016/j.jmb.2016.02.004

- 15 Nys K, Agostinis P, Vermeire S. Autophagy: a new target or an old strategy for the treatment of Crohn's disease? *Nat Rev Gastroenterol Hepatol* 2013;**10**:395–401. doi:10.1038/nrgastro.2013.66
- 16 Dupont N, Lacas-Gervais S, Bertout J, *et al.* Shigella phagocytic vacuolar membrane remnants participate in the cellular response to pathogen invasion and are regulated by autophagy. *Cell Host Microbe* 2009;**6**:137–49. doi:10.1016/j.chom.2009.07.005
- 17 Thurston TLM, Ryzhakov G, Bloor S, *et al.* The TBK1 adaptor and autophagy receptor NDP52 restricts the proliferation of ubiquitin-coated bacteria. *Nat Immunol* 2009;**10**:1215–21. doi:10.1038/ni.1800
- 18 Wild P, Farhan H, McEwan DG, *et al.* Phosphorylation of the autophagy receptor optineurin restricts Salmonella growth. *Science* 2011;**333**:228–33. doi:10.1126/science.1205405
- 19 Johansen T, Lamark T. Selective autophagy mediated by autophagic adapter proteins. *Autophagy* 2011;**7**:279–96.
- 20 Filimonenko M, Isakson P, Finley KD, *et al.* The selective macroautophagic degradation of aggregated proteins requires the PI3P-binding protein Alfy. *Mol Cell* 2010;**38**:265–79. doi:10.1016/j.molcel.2010.04.007
- 21 Kirkin V, Lamark T, Sou Y-S, *et al.* A role for NBR1 in autophagosomal degradation of ubiquitinated substrates. *Mol Cell* 2009;**33**:505–16. doi:10.1016/j.molcel.2009.01.020
- 22 Newman AC, Scholefield CL, Kemp AJ, *et al.* TBK1 kinase addiction in lung cancer cells is mediated via autophagy of Tax1bp1/NDP52 and non-canonical NF- $\kappa$ B signalling. *PloS One* 2012;**7**:e50672. doi:10.1371/journal.pone.0050672
- 23 Smith MD, Harley ME, Kemp AJ, *et al.* CCPG1 Is a Non-canonical Autophagy Cargo Receptor Essential for ER-Phagy and Pancreatic ER Proteostasis. *Dev Cell* 2018;**44**:217–232.e11. doi:10.1016/j.devcel.2017.11.024
- 24 Tschurtschenthaler M, Adolph TE, Ashcroft JW, *et al.* Defective ATG16L1-mediated removal of IRE1 $\alpha$  drives Crohn's disease-like ileitis. *J Exp Med* 2017;**214**:401–22. doi:10.1084/jem.20160791
- 25 Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Cell* 2008;**132**:27–42. doi:10.1016/j.cell.2007.12.018
- 26 Hooper KM, Barlow PG, Stevens C, *et al.* Inflammatory Bowel Disease Drugs: A Focus on Autophagy. *J Crohns Colitis* 2017;**11**:118–27. doi:10.1093/ecco-jcc/jjw127
- 27 Ke P, Shao B-Z, Xu Z-Q, *et al.* Intestinal Autophagy and Its Pharmacological Control in Inflammatory Bowel Disease. *Front Immunol* 2017;**7**. doi:10.3389/fimmu.2016.00695

- 549 28 Hampe J, Franke A, Rosenstiel P, *et al.* A genome-wide association scan of  
1 550 nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in ATG16L1. *Nat*  
2 551 *Genet* 2007;**39**:207–11. doi:10.1038/ng1954
- 5 552 29 Cadwell K, Patel KK, Maloney NS, *et al.* Virus-Plus-Susceptibility Gene Interaction  
6 553 Determines Crohn's Disease Gene Atg16L1 Phenotypes in Intestine. *Cell* 2010;**141**:1135–  
7 554 45. doi:10.1016/j.cell.2010.05.009
- 10 555 30 Cadwell K, Liu J, Brown SL, *et al.* A unique role for autophagy and Atg16L1 in Paneth cells  
11 556 in murine and human intestine. *Nature* 2008;**456**:259–63. doi:10.1038/nature07416
- 13 557 31 Lassen KG, Kuballa P, Conway KL, *et al.* Atg16L1 T300A variant decreases selective  
14 558 autophagy resulting in altered cytokine signaling and decreased antibacterial defense.  
15 559 *Proc Natl Acad Sci U A* 2014;**111**:7741–6. doi:10.1073/pnas.1407001111
- 18 560 32 Kuballa P, Huett A, Rioux JD, *et al.* Impaired autophagy of an intracellular pathogen  
19 561 induced by a Crohn's disease associated ATG16L1 variant. *PLoS One* 2008;**3**:e3391.  
20 562 doi:10.1371/journal.pone.0003391
- 23 563 33 Singh SB, Davis AS, Taylor GA, *et al.* Human IRGM induces autophagy to eliminate  
24 564 intracellular mycobacteria. *Science* 2006;**313**:1438–41. doi:10.1126/science.1129577
- 27 565 34 Brest P, Lapaquette P, Souidi M, *et al.* A synonymous variant in IRGM alters a binding site  
28 566 for miR-196 and causes deregulation of IRGM-dependent xenophagy in Crohn's disease.  
29 567 *Nat Genet* 2011;**43**:242–5. doi:10.1038/ng.762
- 32 568 35 Gardet A, Benita Y, Li C, *et al.* LRRK2 is involved in the IFN-gamma response and host  
33 569 response to pathogens. *J Immunol* 2010;**185**:5577–85. doi:10.4049/jimmunol.1000548
- 36 570 36 Marcuzzi A, Bianco AM, Girardelli M, *et al.* Genetic and functional profiling of Crohn's  
37 571 disease: autophagy mechanism and susceptibility to infectious diseases. *Biomed Res Int*  
38 572 2013;**2013**:297501. doi:10.1155/2013/297501
- 40 573 37 Hugot JP, Chamaillard M, Zouali H, *et al.* Association of NOD2 leucine-rich repeat variants  
41 574 with susceptibility to Crohn's disease. *Nature* 2001;**411**:599–603. doi:10.1038/35079107
- 44 575 38 Rioux JD, Xavier RJ, Taylor KD, *et al.* Genome-wide association study identifies new  
45 576 susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis.  
46 577 *Nat Genet* 2007;**39**:596–604. doi:10.1038/ng2032
- 49 578 39 Weersma RK, Stokkers PCF, van Bodegraven AA, *et al.* Molecular prediction of disease risk  
50 579 and severity in a large Dutch Crohn's disease cohort. *Gut* 2009;**58**:388–95.  
51 580 doi:10.1136/gut.2007.144865
- 54 581 40 Travassos LH, Carneiro LAM, Ramjeet M, *et al.* Nod1 and Nod2 direct autophagy by  
55 582 recruiting ATG16L1 to the plasma membrane at the site of bacterial entry. *Nat Immunol*  
56 583 2010;**11**:55–62. doi:10.1038/ni.1823



- 41 Chauhan S, Mandell MA, Deretic V. IRGM governs the core autophagy machinery to conduct antimicrobial defense. *Mol Cell* 2015;**58**:507–21. doi:10.1016/j.molcel.2015.03.020
- 42 Cooney R, Baker J, Brain O, *et al.* NOD2 stimulation induces autophagy in dendritic cells influencing bacterial handling and antigen presentation. *Nat Med* 2010;**16**:90–7. doi:10.1038/nm.2069
- 43 Homer CR, Richmond AL, Rebert NA, *et al.* ATG16L1 and NOD2 interact in an autophagy-dependent, anti-bacterial pathway implicated in Crohn's disease pathogenesis. *Gastroenterology* 2010;**139**:1630–1641.e2. doi:10.1053/j.gastro.2010.07.006
- 44 Wolfkamp SC, Verseyden C, Vogels EW, *et al.* ATG16L1 and NOD2 polymorphisms enhance phagocytosis in monocytes of Crohn's disease patients., ATG16L1 and NOD2 polymorphisms enhance phagocytosis in monocytes of Crohn's disease patients. *World J Gastroenterol World J Gastroenterol WJG* 2014;**20**, 20:2664, 2664–72. doi:10.3748/wjg.v20.i10.2664, 10.3748/wjg.v20.i10.2664
- 45 Adolph TE, Tomczak MF, Niederreiter L, *et al.* Paneth cells as a site of origin for intestinal inflammation. *Nature* Published Online First: 2 October 2013. doi:10.1038/nature12599
- 46 Deuring JJ, Fuhler GM, Konstantinov SR, *et al.* Genomic ATG16L1 risk allele-restricted Paneth cell ER stress in quiescent Crohn's disease. *Gut* 2014;**63**:1081–91. doi:10.1136/gutjnl-2012-303527
- 47 Kobayashi KS. Nod2-Dependent Regulation of Innate and Adaptive Immunity in the Intestinal Tract. *Science* 2005;**307**:731–4. doi:10.1126/science.1104911
- 48 Wehkamp J, Salzman NH, Porter E, *et al.* Reduced Paneth cell alpha-defensins in ileal Crohn's disease. *Proc Natl Acad Sci U S A* 2005;**102**:18129–34. doi:10.1073/pnas.0505256102
- 49 Liu B, Gulati AS, Cantillana V, *et al.* Irgm1-deficient mice exhibit Paneth cell abnormalities and increased susceptibility to acute intestinal inflammation. *Am J Physiol - Gastrointest Liver Physiol* 2013;**305**:G573–84. doi:10.1152/ajpgi.00071.2013
- 50 Zhang Q, Pan Y, Yan R, *et al.* Commensal bacteria direct selective cargo sorting to promote symbiosis. *Nat Immunol* 2015;**16**:918–26. doi:10.1038/ni.3233
- 51 Saitoh T, Fujita N, Jang MH, *et al.* Loss of the autophagy protein Atg16L1 enhances endotoxin-induced IL-1 $\beta$  production. *Nature* 2008;**456**:264–8. doi:10.1038/nature07383
- 52 Zhang H, Zheng L, McGovern DPB, *et al.* Myeloid ATG16L1 Facilitates Host-Bacteria Interactions in Maintaining Intestinal Homeostasis. *J Immunol Baltim Md 1950* 2017;**198**:2133–46. doi:10.4049/jimmunol.1601293
- 53 Mizoguchi A, Bhan AK. Immunobiology of B Cells in Inflammatory Bowel Disease. In: *Crohn's Disease and Ulcerative Colitis*. Springer, Boston, MA 2012. 161–8. doi:10.1007/978-1-4614-0998-4\_12

- 54 Maeda S, Hsu L-C, Liu H, *et al.* Nod2 mutation in Crohn's disease potentiates NF-kappaB activity and IL-1beta processing. *Science* 2005;**307**:734–8. doi:10.1126/science.1103685
- 55 Liu Z, Lee J, Krummey S, *et al.* The kinase LRRK2 is a regulator of the transcription factor NFAT that modulates the severity of inflammatory bowel disease. *Nat Immunol* 2011;**12**:1063–70. doi:10.1038/ni.2113
- 56 Sano R, Reed JC. ER stress-induced cell death mechanisms. *Biochim Biophys Acta* 2013;**1833**. doi:10.1016/j.bbamcr.2013.06.028
- 57 Cao SS. Endoplasmic Reticulum Stress and Unfolded Protein Response in Inflammatory Bowel Disease: *Inflamm Bowel Dis* 2015;**21**:636–44. doi:10.1097/MIB.0000000000000238
- 58 Guan B-J, Krokowski D, Majumder M, *et al.* Translational control during endoplasmic reticulum stress beyond phosphorylation of the translation initiation factor eIF2 $\alpha$ . *J Biol Chem* 2014;**289**:12593–611. doi:10.1074/jbc.M113.543215
- 59 Vattem KM, Wek RC. Reinitiation involving upstream ORFs regulates ATF4 mRNA translation in mammalian cells. *Proc Natl Acad Sci U S A* 2004;**101**:11269–74. doi:10.1073/pnas.0400541101
- 60 Harding HP, Zhang Y, Bertolotti A, *et al.* Perk Is Essential for Translational Regulation and Cell Survival during the Unfolded Protein Response. *Mol Cell* 2000;**5**:897–904. doi:10.1016/S1097-2765(00)80330-5
- 61 Nishitoh H. CHOP is a multifunctional transcription factor in the ER stress response. *J Biochem (Tokyo)* 2012;**151**:217–9. doi:10.1093/jb/mvr143
- 62 Wang XZ, Harding HP, Zhang Y, *et al.* Cloning of mammalian Ire1 reveals diversity in the ER stress responses. *EMBO J* 1998;**17**:5708–17. doi:10.1093/emboj/17.19.5708
- 63 Shamu CE, Walter P. Oligomerization and phosphorylation of the Ire1p kinase during intracellular signaling from the endoplasmic reticulum to the nucleus. *EMBO J* 1996;**15**:3028–39.
- 64 Tirasophon W, Lee K, Callaghan B, *et al.* The endoribonuclease activity of mammalian IRE1 autoregulates its mRNA and is required for the unfolded protein response. *Genes Dev* 2000;**14**:2725–36.
- 65 Calfon M, Zeng H, Urano F, *et al.* IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. *Nature* 2002;**415**:92–6. doi:10.1038/415092a
- 66 Lee A-H, Iwakoshi NN, Glimcher LH. XBP-1 Regulates a Subset of Endoplasmic Reticulum Resident Chaperone Genes in the Unfolded Protein Response. *Mol Cell Biol* 2003;**23**:7448–59. doi:10.1128/MCB.23.21.7448-7459.2003

- 67 Lee K, Tirasophon W, Shen X, *et al.* IRE1-mediated unconventional mRNA splicing and S2P-mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein response. *Genes Dev* 2002;**16**:452–66. doi:10.1101/gad.964702
- 68 Yoshida H, Matsui T, Yamamoto A, *et al.* XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* 2001;**107**:881–91.
- 69 Hollien J. Decay of Endoplasmic Reticulum-Localized mRNAs During the Unfolded Protein Response. *Science* 2006;**313**:104–7. doi:10.1126/science.1129631
- 70 Shen J, Chen X, Hendershot L, *et al.* ER stress regulation of ATF6 localization by dissociation of BiP/GRP78 binding and unmasking of Golgi localization signals. *Dev Cell* 2002;**3**:99–111.
- 71 Haze K, Yoshida H, Yanagi H, *et al.* Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress. *Mol Biol Cell* 1999;**10**:3787–99.
- 72 Li M, Baumeister P, Roy B, *et al.* ATF6 as a transcription activator of the endoplasmic reticulum stress element: thapsigargin stress-induced changes and synergistic interactions with NF-Y and YY1. *Mol Cell Biol* 2000;**20**:5096–106.
- 73 Ye J, Rawson RB, Komuro R, *et al.* ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs. *Mol Cell* 2000;**6**:1355–64.
- 74 Hirsch I, Weiwad M, Prell E, *et al.* ERp29 deficiency affects sensitivity to apoptosis via impairment of the ATF6-CHOP pathway of stress response. *Apoptosis Int J Program Cell Death* 2014;**19**:801–15. doi:10.1007/s10495-013-0961-0
- 75 McGuckin MA, Eri RD, Das I, *et al.* ER stress and the unfolded protein response in intestinal inflammation. *Am J Physiol - Gastrointest Liver Physiol* 2010;**298**:G820–32. doi:10.1152/ajpgi.00063.2010
- 76 Deuring JJ, de Haar C, Koelewijn CL, *et al.* Absence of ABCG2-mediated mucosal detoxification in patients with active inflammatory bowel disease is due to impeded protein folding. *Biochem J* 2012;**441**:87–93. doi:10.1042/BJ20111281
- 77 Rolhion N, Barnich N, Bringer M-A, *et al.* Abnormally expressed ER stress response chaperone Gp96 in CD favours adherent-invasive Escherichia coli invasion. *Gut* 2010;**59**:1355–62. doi:10.1136/gut.2010.207456
- 78 Shkoda A, Ruiz PA, Daniel H, *et al.* Interleukin-10 Blocked Endoplasmic Reticulum Stress in Intestinal Epithelial Cells: Impact on Chronic Inflammation. *Gastroenterology* 2007;**132**:190–207. doi:10.1053/j.gastro.2006.10.030
- 79 Niederreiter L, Fritz TMJ, Adolph TE, *et al.* ER stress transcription factor Xbp1 suppresses intestinal tumorigenesis and directs intestinal stem cells. *J Exp Med* 2013;**210**:2041–56. doi:10.1084/jem.20122341

- 692 80 Martinon F, Chen X, Lee A-H, *et al.* TLR activation of the transcription factor XBP1  
693 regulates innate immune responses in macrophages. *Nat Immunol* 2010;**11**:411–8.  
694 doi:10.1038/ni.1857
- 695 81 Bertolotti A, Wang X, Novoa I, *et al.* Increased sensitivity to dextran sodium sulfate colitis  
696 in IRE1 $\beta$ -deficient mice. *J Clin Invest* 2001;**107**:585–93.
- 697 82 Tsuru A, Fujimoto N, Takahashi S, *et al.* Negative feedback by IRE1 $\beta$  optimizes mucin  
698 production in goblet cells. *Proc Natl Acad Sci U S A* 2013;**110**:2864–9.  
699 doi:10.1073/pnas.1212484110
- 700 83 Cao SS, Wang M, Harrington JC, *et al.* Phosphorylation of eIF2 $\alpha$  Is Dispensable for  
701 Differentiation but Required at a Posttranscriptional Level for Paneth Cell Function and  
702 Intestinal Homeostasis in Mice: *Inflamm Bowel Dis* 2014;**20**:712–22.  
703 doi:10.1097/MIB.0000000000000010
- 704 84 Cao SS, Zimmermann EM, Chuang B, *et al.* The Unfolded Protein Response and Chemical  
705 Chaperones Reduce Protein Misfolding and Colitis in Mice. *Gastroenterology*  
706 2013;**144**:989–1000.e6. doi:10.1053/j.gastro.2013.01.023
- 707 85 Brandl K, Rutschmann S, Li X, *et al.* Enhanced sensitivity to DSS colitis caused by a  
708 hypomorphic Mbtsp1 mutation disrupting the ATF6-driven unfolded protein response.  
709 *Proc Natl Acad Sci U S A* 2009;**106**:3300–5. doi:10.1073/pnas.0813036106
- 710 86 Moehle C, Ackermann N, Langmann T, *et al.* Aberrant intestinal expression and allelic  
711 variants of mucin genes associated with inflammatory bowel disease. *J Mol Med Berl Ger*  
712 2006;**84**:1055–66. doi:10.1007/s00109-006-0100-2
- 713 87 Eri RD, Adams RJ, Tran TV, *et al.* An intestinal epithelial defect conferring ER stress results  
714 in inflammation involving both innate and adaptive immunity. *Mucosal Immunol*  
715 2011;**4**:354–64. doi:10.1038/mi.2010.74
- 716 88 Asada R, Saito A, Kawasaki N, *et al.* The Endoplasmic Reticulum Stress Transducer OASIS  
717 Is involved in the Terminal Differentiation of Goblet Cells in the Large Intestine. *J Biol*  
718 *Chem* 2012;**287**:8144–53. doi:10.1074/jbc.M111.332593
- 719 89 Hino K, Saito A, Asada R, *et al.* Increased Susceptibility to Dextran Sulfate Sodium-Induced  
720 Colitis in the Endoplasmic Reticulum Stress Transducer OASIS Deficient Mice. *PLoS ONE*  
721 2014;**9**. doi:10.1371/journal.pone.0088048
- 722 90 Zhao F, Edwards R, Dizon D, *et al.* Disruption of Paneth and goblet cell homeostasis and  
723 increased endoplasmic reticulum stress in Agr2 $^{-/-}$  mice. *Dev Biol* 2010;**338**:270–9.  
724 doi:10.1016/j.ydbio.2009.12.008
- 725 91 Shimodaira Y, Takahashi S, Kinouchi Y, *et al.* Modulation of endoplasmic reticulum (ER)  
726 stress-induced autophagy by C/EBP homologous protein (CHOP) and inositol-requiring  
727 enzyme 1 $\alpha$  (IRE1 $\alpha$ ) in human colon cancer cells. *Biochem Biophys Res Commun*  
728 2014;**445**:524–33. doi:10.1016/j.bbrc.2014.02.054

- 92 Margariti A, Li H, Chen T, *et al.* XBP1 mRNA Splicing Triggers an Autophagic Response in Endothelial Cells through BECLIN-1 Transcriptional Activation. *J Biol Chem* 2013;**288**:859–72. doi:10.1074/jbc.M112.412783
- 93 Hetz C, Thielen P, Matus S, *et al.* XBP-1 deficiency in the nervous system protects against amyotrophic lateral sclerosis by increasing autophagy. *Genes Dev* 2009;**23**:2294–306. doi:10.1101/gad.1830709
- 94 Avivar-Valderas A, Bobrovnikova-Marjon E, Diehl JA, *et al.* Regulation of autophagy during ECM detachment is linked to a selective inhibition of mTORC1 by PERK. *Oncogene* 2013;**32**:4932–40. doi:10.1038/onc.2012.512
- 95 Ji G, Yu N, Xue X, *et al.* PERK-mediated Autophagy in Osteosarcoma Cells Resists ER Stress-induced Cell Apoptosis. *Int J Biol Sci* 2015;**11**:803–12. doi:10.7150/ijbs.11100
- 96 Jia X-E, Ma K, Xu T, *et al.* Mutation of kri1l causes definitive hematopoiesis failure via PERK-dependent excessive autophagy induction. *Cell Res* 2015;**25**:946.
- 97 Kouroku Y, Fujita E, Tanida I, *et al.* ER stress (PERK/eIF2 [alpha] phosphorylation) mediates the polyglutamine-induced LC3 conversion, an essential step for autophagy formation. *Cell Death Differ* 2007;**14**:230.
- 98 Moon H, Kim B, Gwak H, *et al.* Autophagy and protein kinase RNA-like endoplasmic reticulum kinase (PERK)/eukaryotic initiation factor 2 alpha kinase (eIF2 $\alpha$ ) pathway protect ovarian cancer cells from metformin-induced apoptosis: THE EFFECT OF METFORMIN ON AUTOPHAGY AND PERK. *Mol Carcinog* 2016;**55**:346–56. doi:10.1002/mc.22284
- 99 Zhao C, Yin S, Dong Y, *et al.* Autophagy-dependent EIF2AK3 activation compromises ursolic acid-induced apoptosis through upregulation of MCL1 in MCF-7 human breast cancer cells. *Autophagy* 2013;**9**:196–207. doi:10.4161/auto.22805
- 100 Laplante M, Sabatini DM. mTOR Signaling in Growth Control and Disease. *Cell* 2012;**149**:274–93. doi:10.1016/j.cell.2012.03.017
- 101 B'chir W, Maurin A-C, Carraro V, *et al.* The eIF2 $\alpha$ /ATF4 pathway is essential for stress-induced autophagy gene expression. *Nucleic Acids Res* 2013;**41**:7683–99. doi:10.1093/nar/gkt563
- 102 Avivar-Valderas A, Salas E, Bobrovnikova-Marjon E, *et al.* PERK Integrates Autophagy and Oxidative Stress Responses To Promote Survival during Extracellular Matrix Detachment. *Mol Cell Biol* 2011;**31**:3616–29. doi:10.1128/MCB.05164-11
- 103 Rouschop KMA, van den Beucken T, Dubois L, *et al.* The unfolded protein response protects human tumor cells during hypoxia through regulation of the autophagy genes MAP1LC3B and ATG5. *J Clin Invest* 2010;**120**:127–41. doi:10.1172/JCI40027
- 104 Rzymiski T, Milani M, Pike L, *et al.* Regulation of autophagy by ATF4 in response to severe hypoxia. *Oncogene* 2010;**29**:4424–35. doi:10.1038/onc.2010.191

- 105 Gade P, Manjegowda SB, Nallar SC, *et al.* Regulation of the Death-Associated Protein Kinase 1 Expression and Autophagy via ATF6 Requires Apoptosis Signal-Regulating Kinase 1. *Mol Cell Biol* 2014;**34**:4033–48. doi:10.1128/MCB.00397-14
- 106 Gade P, Ramachandran G, Maachani UB, *et al.* An IFN- $\gamma$ -stimulated ATF6–C/EBP- $\beta$ -signaling pathway critical for the expression of Death Associated Protein Kinase 1 and induction of autophagy. *Proc Natl Acad Sci U S A* 2012;**109**:10316–21. doi:10.1073/pnas.1119273109
- 107 Lopes F, Keita ÅV, Saxena A, *et al.* ER-stress mobilization of death-associated protein kinase-1-dependent xenophagy counteracts mitochondria stress-induced epithelial barrier dysfunction. *J Biol Chem* 2018;**293**:3073–87. doi:10.1074/jbc.RA117.000809
- 108 Appenzeller-Herzog C, Hall MN. Bidirectional crosstalk between endoplasmic reticulum stress and mTOR signaling. *Trends Cell Biol* 2012;**22**:274–82. doi:10.1016/j.tcb.2012.02.006
- 109 Yamazaki H, Hiramatsu N, Hayakawa K, *et al.* Activation of the Akt-NF-kappaB pathway by subtilase cytotoxin through the ATF6 branch of the unfolded protein response. *J Immunol Baltim Md 1950* 2009;**183**:1480–7. doi:10.4049/jimmunol.0900017
- 110 Hart LS, Cunningham JT, Datta T, *et al.* ER stress-mediated autophagy promotes Myc-dependent transformation and tumor growth. *J Clin Invest* 2012;**122**:4621–34. doi:10.1172/JCI62973
- 111 Li J, Ni M, Lee B, *et al.* The unfolded protein response regulator GRP78/BiP is required for endoplasmic reticulum integrity and stress-induced autophagy in mammalian cells. *Cell Death Differ* 2008;**15**:1460–71. doi:10.1038/cdd.2008.81
- 112 Ogata M, Hino S, Saito A, *et al.* Autophagy is activated for cell survival after endoplasmic reticulum stress. *Mol Cell Biol* 2006;**26**:9220–31. doi:10.1128/mcb.01453-06
- 113 Wang W, Kang H, Zhao Y, *et al.* Targeting autophagy sensitizes BRAF-mutant thyroid cancer to vemurafenib. *J Clin Endocrinol Metab* 2016;**jc.2016-1999**. doi:10.1210/jc.2016-1999
- 114 Keestra-Gounder AM, Byndloss MX, Seyffert N, *et al.* NOD1 and NOD2 signalling links ER stress with inflammation. *Nature* 2016;**532**:394–7. doi:10.1038/nature17631
- 115 Kaneko M, Niinuma Y, Nomura Y. Activation Signal of Nuclear Factor- $\kappa$ B in Response to Endoplasmic Reticulum Stress is Transduced via IRE1 and Tumor Necrosis Factor Receptor-Associated Factor 2. *Biol Pharm Bull* 2003;**26**:931–5. doi:10.1248/bpb.26.931
- 116 Urano F, Wang X, Bertolotti A, *et al.* Coupling of Stress in the ER to Activation of JNK Protein Kinases by Transmembrane Protein Kinase IRE1. *Science* 2000;**287**:664–6. doi:10.1126/science.287.5453.664

- 117 Castillo K, Rojas-Rivera D, Lisbona F, *et al.* BAX inhibitor-1 regulates autophagy by controlling the IRE1 $\alpha$  branch of the unfolded protein response. *EMBO J* 2011;**30**:4465–78. doi:10.1038/emboj.2011.318
- 118 Ding W-X, Ni H-M, Gao W, *et al.* Linking of Autophagy to Ubiquitin-Proteasome System Is Important for the Regulation of Endoplasmic Reticulum Stress and Cell Viability. *Am J Pathol* 2007;**171**:513–24. doi:10.2353/ajpath.2007.070188
- 119 Moretti J, Roy S, Bozec D, *et al.* STING Senses Microbial Viability to Orchestrate Stress-Mediated Autophagy of the Endoplasmic Reticulum. *Cell* Published Online First: October 2017. doi:10.1016/j.cell.2017.09.034
- 120 NHS CB. 2013/14 NHS Standard Contract for Colorectal: Complex Inflammatory Bowel Disease (Adult). 2013.
- 121 Bassi A, Dodd S, Williamson P, *et al.* Cost of illness of inflammatory bowel disease in the UK: a single centre retrospective study. *Gut* 2004;**53**:1471–8. doi:10.1136/gut.2004.041616
- 122 Bernstein CN, Loftus EV Jr, Ng SC, *et al.* Hospitalisations and surgery in Crohn's disease. *Gut* 2012;**61**:622–9. doi:10.1136/gutjnl-2011-301397
- 123 Matsuda C, Ito T, Song J, *et al.* Therapeutic effect of a new immunosuppressive agent, everolimus, on interleukin-10 gene-deficient mice with colitis. *Clin Exp Immunol* 2007;**148**:348–59. doi:10.1111/j.1365-2249.2007.03345.x
- 124 Massey DC, Bredin F, Parkes M. Use of sirolimus (rapamycin) to treat refractory Crohn's disease. *Gut* 2008;**57**:1294–6. doi:10.1136/gut.2008.157297
- 125 Dumortier J, Lapalus M-G, Guillaud O, *et al.* Everolimus for refractory Crohn's disease: A case report: *Inflamm Bowel Dis* 2008;**14**:874–7. doi:10.1002/ibd.20395
- 126 Mutalib M, Borrelli O, Blackstock S, *et al.* The use of sirolimus (rapamycin) in the management of refractory inflammatory bowel disease in children. *J Crohns Colitis* 2014;**8**:1730–4. doi:10.1016/j.crohns.2014.08.014
- 127 Reinisch W, Panés J, Lémann M, *et al.* A multicenter, randomized, double-blind trial of everolimus versus azathioprine and placebo to maintain steroid-induced remission in patients with moderate-to-severe active Crohn's disease. *Am J Gastroenterol* 2008;**103**:2284–92. doi:10.1111/j.1572-0241.2008.02024.x
- 128 Ha J, Kim J. Novel pharmacological modulators of autophagy: an updated patent review (2012-2015). *Expert Opin Ther Pat* 2016;**26**:1273–89. doi:10.1080/13543776.2016.1217996
- 129 Galluzzi L, Bravo-San Pedro JM, Levine B, *et al.* Pharmacological modulation of autophagy: therapeutic potential and persisting obstacles. *Nat Rev Drug Discov* 2017;**16**:487–511. doi:10.1038/nrd.2017.22

- 130 Rauthan M, Pilon M. A chemical screen to identify inducers of the mitochondrial unfolded protein response in *C. elegans*. *Worm* 2015;**4**. doi:10.1080/21624054.2015.1096490
- 131 Boyce M, Bryant KF, Jousse C, *et al*. A selective inhibitor of eIF2alpha dephosphorylation protects cells from ER stress. *Science* 2005;**307**:935–9. doi:10.1126/science.1101902
- 132 Okazaki T, Nishio A, Takeo M, *et al*. Inhibition of the dephosphorylation of eukaryotic initiation factor 2 $\alpha$  ameliorates murine experimental colitis. *Digestion* 2014;**90**:167–78. doi:10.1159/000366414
- 133 Crespo I, San-Miguel B, Prause C, *et al*. Glutamine treatment attenuates endoplasmic reticulum stress and apoptosis in TNBS-induced colitis. *PloS One* 2012;**7**:e50407. doi:10.1371/journal.pone.0050407
- 134 Yang L, Sha H, Davisson RL, *et al*. Phenformin activates the unfolded protein response in an AMP-activated protein kinase (AMPK)-dependent manner. *J Biol Chem* 2013;**288**:13631–8. doi:10.1074/jbc.M113.462762
- 135 Kim H, Moon SY, Kim J-S, *et al*. Activation of AMP-activated protein kinase inhibits ER stress and renal fibrosis. *Am J Physiol Renal Physiol* 2015;**308**:F226–236. doi:10.1152/ajprenal.00495.2014
- 136 Hwang H-J, Jung TW, Ryu JY, *et al*. Dipeptidyl peptidase-IV inhibitor (gemigliptin) inhibits tunicamycin-induced endoplasmic reticulum stress, apoptosis and inflammation in H9c2 cardiomyocytes. *Mol Cell Endocrinol* 2014;**392**:1–7. doi:10.1016/j.mce.2014.04.017
- 137 Yusta B, Baggio LL, Estall JL, *et al*. GLP-1 receptor activation improves beta cell function and survival following induction of endoplasmic reticulum stress. *Cell Metab* 2006;**4**:391–406. doi:10.1016/j.cmet.2006.10.001
- 138 Engin F, Hotamisligil GS. Restoring endoplasmic reticulum function by chemical chaperones: an emerging therapeutic approach for metabolic diseases. *Diabetes Obes Metab* 2010;**12 Suppl 2**:108–15. doi:10.1111/j.1463-1326.2010.01282.x
- 139 Allaire JM, Crowley SM, Law HT, *et al*. The Intestinal Epithelium: Central Coordinator of Mucosal Immunity. *Trends Immunol* 2018;**39**:677–96. doi:10.1016/j.it.2018.04.002
- 140 Boyapati RK, Rossi AG, Satsangi J, *et al*. Gut mucosal DAMPs in IBD: from mechanisms to therapeutic implications. *Mucosal Immunol* 2016;**9**:567–82. doi:10.1038/mi.2016.14
- 141 Boyapati RK, Dorward DA, Tamborska A, *et al*. Mitochondrial DNA Is a Pro-Inflammatory Damage-Associated Molecular Pattern Released During Active IBD. *Inflamm Bowel Dis* Published Online First: 1 May 2018. doi:10.1093/ibd/izy095
- 142 Odenwald MA, Turner JR. The intestinal epithelial barrier: a therapeutic target? *Nat Rev Gastroenterol Hepatol* 2017;**14**:9–21. doi:10.1038/nrgastro.2016.169



- 143 Nishino K, Nishida A, Inoue R, *et al.* Analysis of endoscopic brush samples identified mucosa-associated dysbiosis in inflammatory bowel disease. *J Gastroenterol* 2018;**53**:95–106. doi:10.1007/s00535-017-1384-4
- 144 Wildenberg ME, Koelink PJ, Diederens K, *et al.* The ATG16L1 risk allele associated with Crohn's disease results in a Rac1-dependent defect in dendritic cell migration that is corrected by thiopurines. *Mucosal Immunol* 2017;**10**:352–60. doi:10.1038/mi.2016.65
- 145 Wildenberg M, Levin A, Vos C, *et al.* P668 ATG16L1 genotype is associated with response to anti-TNF in vitro. *J Crohns Colitis* 2013;**7**:S279. doi:10.1016/s1873-9946(13)60689-3
- 146 Levin AD, Koelink PJ, Bloemendaal FM, *et al.* Autophagy Contributes to the Induction of Anti-TNF Induced Macrophages. *J Crohns Colitis* 2016;**10**:323–9. doi:10.1093/ecco-jcc/jjv174
- 147 Vos ACW, Wildenberg ME, Arijis I, *et al.* Regulatory macrophages induced by infliximab are involved in healing in vivo and in vitro. *Inflamm Bowel Dis* 2012;**18**:401–8. doi:10.1002/ibd.21818
- 148 Niess JH, Klaus J, Stephani J, *et al.* NOD2 polymorphism predicts response to treatment in Crohn's disease--first steps to a personalized therapy. *Dig Sci* 2012;**57**:879–86. doi:10.1007/s10620-011-1977-3
- 149 Chouchana L, Fernández-Ramos AA, Dumont F, *et al.* Molecular insight into thiopurine resistance: transcriptomic signature in lymphoblastoid cell lines. *Genome Med* 2015;**7**:37. doi:10.1186/s13073-015-0150-6
- 150 Ventham NT, Kennedy NA, Nimmo ER, *et al.* Beyond gene discovery in inflammatory bowel disease: the emerging role of epigenetics. *Gastroenterology* 2013;**145**:293–308. doi:10.1053/j.gastro.2013.05.050
- 151 Kalla R, Ventham NT, Kennedy NA, *et al.* MicroRNAs: new players in IBD. *Gut* 2015;**64**:504–17. doi:10.1136/gutjnl-2014-307891
- 152 Stevens TW, Matheeuwsen M, Lönnkvist MH, *et al.* Systematic review: predictive biomarkers of therapeutic response in inflammatory bowel disease-Personalised medicine in its infancy. *Aliment Pharmacol Ther* Published Online First: 30 October 2018. doi:10.1111/apt.15033
- 153 Pascal V, Pozuelo M, Borruel N, *et al.* A microbial signature for Crohn's disease. *Gut* 2017;;gutjnl-2016-313235. doi:10.1136/gutjnl-2016-313235